

SAMPLE OUTLINE OF LETTER OF APPEAL CHANGE OF TREATMENT

[Date]

[Health plan name]

ATTN: [Department]

[Medical/Pharmacy Director Name (if available)]

[Health plan address]

[City, State, ZIP code]

[Patient's Name]

[Patient's plan-specific member ID]

[Date of birth]

[Case number]

[Dates of service]

Re: Appeal of Denial for KESIMPTA® (ofatumumab)

Dear [Medical/Pharmacy Director Name],

This letter of [level of appeal] is a formal appeal of your coverage decision for KESIMPTA for the patient referenced above. The reason(s) for the denial were [List reason(s) for the denial]. I request that [Insurance name]'s denial decision be reversed and coverage approved for KESIMPTA as it is medically necessary to treat the diagnosis of relapsing multiple sclerosis (RMS) [(ICD-10 code)]. At this time, the appropriate treatment plan is to discontinue [Current Drug] and to prescribe KESIMPTA.

I have been treating [Patient's Name], [a/an] [age]-year-old [male/female], since [Date] to manage RMS [(ICD-10 code)]. [He/She] has been on [Drug name] since [Date].

[Include relevant medical information to support your reason for discontinuing the current medication and prescribing KESIMPTA. If applicable, include evidence that the patient's RMS symptoms and disabilities have been progressing despite his/her current therapies. Additional information may include:

- Supporting information as requested by the plan in its denial letter
- Clinical attributes of KESIMPTA and relevance to patient]

History of previous multiple sclerosis (MS) therapies: _____

Reasons for discontinuation of previous therapies: _____

Duration of previous therapies: _____

Based on the patient's condition and medical history, as well as my experience in treating patients with RMS/MS [(ICD-10 code)], I believe treatment with KESIMPTA is appropriate and medically necessary.

Included with this letter of appeal for approval to change to KESIMPTA are relevant supporting medical documentation, including a letter of medical necessity, clinical trial information, and the US Food and Drug Administration (FDA) approval letter. I have also attached the original denial letter. [Summarize reasons for the patient to convert to KESIMPTA].

If you have any further questions about this matter, please feel free to contact me at [physician phone number] or via email at [physician email]. Thank you for your time and consideration.

Sincerely,

[Physician's signature]

Enclosures

[List and attach additional documents, which may include a denial letter, Letter of Medical Necessity, Prescribing Information, clinical notes/medical records, FDA approval letter, or clinical practice guidelines.]

This letter is provided as an example and is meant for educational purposes only. Novartis cannot guarantee insurance coverage or reimbursement. Coverage and reimbursement may vary significantly by payer, plan, patient, and setting of care. It is the sole responsibility of the health care provider to include the proper information and ensure the accuracy of all statements used in seeking coverage and reimbursement for an individual patient.

Click [here](#) for full Prescribing Information, including Medication Guide.

Indication and Important Safety Information

INDICATION

KESIMPTA is indicated for the treatment of relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

IMPORTANT SAFETY INFORMATION

Contraindications

KESIMPTA is contraindicated in patients with active hepatitis B virus (HBV) infection, or history of hypersensitivity to ofatumumab, or life-threatening injection-related reaction to KESIMPTA. Hypersensitivity reactions have included anaphylaxis and angioedema.

Warnings and Precautions

Infections

Serious, including life-threatening or fatal, bacterial, fungal, and new or reactivated viral infections have been observed during and following completion of treatment with anti-CD20 B-cell depleting therapies. The overall rate of infections and serious infections in KESIMPTA-treated patients was similar to triflunomide-treated patients (51.6% vs 52.7%, and 2.5% vs 1.8%, respectively). The most common infections reported by KESIMPTA-treated patients in relapsing MS (RMS) trials included upper respiratory tract infection (39%) and urinary tract infection (10%). Delay KESIMPTA administration in patients with an active infection until resolved.

Consider the potential increased immunosuppressive effects when initiating KESIMPTA after an immunosuppressive therapy or initiating an immunosuppressive therapy after KESIMPTA.

Hepatitis B Virus

Reactivation: No reports of HBV reactivation in patients with MS treated with KESIMPTA. However, HBV reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, has occurred in patients treated with ofatumumab at higher intravenous doses for chronic lymphocytic leukemia (CLL) than the recommended dose in MS and in patients treated with other anti-CD20 antibodies.

Infection: KESIMPTA is contraindicated in patients with active hepatitis B disease. Fatal infections caused by HBV in patients who have not been previously infected have occurred in patients treated with ofatumumab at higher intravenous doses for CLL than the recommended dose in MS. Perform HBV screening in all patients before initiation of KESIMPTA. Patients who are negative for HBsAg and positive for HB core antibody [HBcAb+] or are carriers of HBV [HBsAg+], should consult liver disease experts before starting and during KESIMPTA treatment.

Progressive Multifocal Leukoencephalopathy

No cases of progressive multifocal leukoencephalopathy (PML) have been reported for KESIMPTA in RMS clinical studies; however, PML resulting in death has occurred in patients being treated with ofatumumab at higher intravenous doses for CLL than the recommended dose in MS. In addition, JC virus infection resulting in PML has also been observed in patients treated with other anti-CD20 antibodies and other MS therapies. If PML is suspected, withhold KESIMPTA and perform an appropriate diagnostic evaluation. If PML is confirmed, KESIMPTA should be discontinued.

Vaccinations

Administer all immunizations according to immunization guidelines: for live or live-attenuated vaccines at least 4 weeks and, whenever possible at least 2 weeks prior to starting KESIMPTA for inactivated vaccines. The safety of immunization with live or live-attenuated vaccines following KESIMPTA therapy has not been studied. Vaccination with live or live-attenuated vaccines is not recommended during treatment and after discontinuation until B-cell repletion.

Vaccination of Infants Born to Mothers Treated with KESIMPTA During Pregnancy

For infants whose mother was treated with KESIMPTA during pregnancy, assess B-cell

counts prior to administration of live or live-attenuated vaccines. If the B-cell count has not recovered in the infant, do not administer the vaccine as having depleted B-cells may pose an increased risk in these infants.

Injection-Related Reactions and Hypersensitivity Reactions

KESIMPTA can result in systemic injection-related reactions and hypersensitivity reactions, which may be serious or life-threatening. Injection-related reactions with systemic symptoms occurred most commonly within 24 hours of the first injection, but were also observed with later injections. There were no life-threatening injection reactions in RMS clinical studies.

In the post-marketing setting, additional systemic injection-related reactions and hypersensitivity reactions have been reported, including anaphylaxis, angioedema, pruritus, rash, urticaria, erythema, bronchospasm, throat irritation, oropharyngeal pain, dyspnea, pharyngeal or laryngeal edema, flushing, hypotension, dizziness, nausea, and tachycardia. Most cases were not serious and occurred with the first injection. Symptoms of systemic injection-related reactions may be clinically indistinguishable from acute hypersensitivity reactions.

The first injection of KESIMPTA should be performed under the guidance of an appropriately trained health care professional. If systemic injection-related reactions occur, initiate appropriate therapy. Patients who experience symptoms of systemic injection-related reactions or hypersensitivity reactions with KESIMPTA should be instructed to seek immediate medical attention. If local injection-related reactions occur, symptomatic treatment is recommended.

Reduction in Immunoglobulins

As expected with any B-cell depleting therapy, decreased immunoglobulin levels were observed. Monitor the levels of quantitative serum immunoglobulins during treatment, especially in patients with opportunistic or recurrent infections and after discontinuation of therapy until B-cell repletion. Consider discontinuing KESIMPTA therapy if a patient with low immunoglobulins develops a serious opportunistic infection or recurrent infections, or if prolonged hypogammaglobulinemia requires treatment with intravenous immunoglobulins.

Liver Injury

Clinically significant liver injury, without findings of viral hepatitis, has been reported in the post-marketing setting. Signs of liver injury have occurred weeks to months after administration. Patients treated with KESIMPTA found to have an alanine aminotransferase or aspartate aminotransferase greater than 3 times the upper limit of normal (ULN) with serum total bilirubin greater than 2 times the ULN are potentially at risk for severe drug-induced liver injury.

Obtain liver function tests prior to initiating treatment. Monitor for signs and symptoms of hepatic injury during treatment, including new or worsening fatigue, anorexia, nausea, vomiting, right upper abdominal discomfort, dark urine, or jaundice. If symptoms of liver injury are reported, measure serum aminotransferases, alkaline phosphatase, and bilirubin levels. Discontinue KESIMPTA if liver injury is present and an alternative etiology is not identified.

Fetal Risk

Based on animal data, KESIMPTA can cause fetal harm due to B-cell lymphopenia and reduce antibody response in offspring exposed to KESIMPTA in utero. Transient peripheral B-cell depletion and lymphocytopenia have been reported in infants born to mothers exposed to other anti-CD20 B-cell depleting antibodies during pregnancy. Advise females of reproductive potential to use effective contraception while receiving KESIMPTA and for at least 6 months after the last dose.

Most Common Adverse Reactions

Most common adverse reactions (>10%) are upper respiratory tract infection, headache, injection-related reactions, and local injection-site reactions.