



Start Form

Initiate treatment and patient support
with a 1-page form

INDICATION

KESIMPTA is indicated for the treatment of relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

IMPORTANT SAFETY INFORMATION

Contraindications: KESIMPTA is contraindicated in patients with active hepatitis B virus (HBV) infection, or history of hypersensitivity to ofatumumab, or life-threatening injection-related reaction to KESIMPTA. Hypersensitivity reactions have included anaphylaxis and angioedema.

WARNINGS AND PRECAUTIONS

Infections: Serious, including life-threatening or fatal, bacterial, fungal, and new or reactivated viral infections have been observed during and following completion of treatment with anti-CD20 B-cell depleting therapies. The overall rate of infections and serious infections in KESIMPTA-treated patients was similar to teriflunomide-treated patients (51.6% vs 52.7%, and 2.5% vs 1.8%, respectively). The most common infections reported by KESIMPTA-treated patients in relapsing MS (RMS) trials included upper respiratory tract infection (39%) and urinary tract infection (10%). Delay KESIMPTA administration in patients with an active infection until resolved.

Consider the potential increased immunosuppressive effects when initiating KESIMPTA after an immunosuppressive therapy or initiating an immunosuppressive therapy after KESIMPTA.

Please see additional Important Safety Information on [the last page](#).

Please see full Prescribing Information, including Medication Guide [here](#).

Get started with the Start Form

It's pretty straightforward, but we highlighted a few things to keep in mind. We're only asking for essential information to make it easier on you. Fill it out completely to help make getting started **faster for your patients.**

Get patient and/or guardian consent.

Check this box to sign patients up for the \$0 Access Card, which includes access to the Bridge and Copay Programs.

Don't skip the prescription insurance info! We need it to verify all your patient's benefits.

Sign up for Bridge to Commercial Coverage and the Novartis Patient Assistance Foundation (NPAF) for access to loading and/or maintenance doses.

Make sure to provide a prescriber signature, too!

KESIMPTA®
Prescription Start Form

Send Fax 1-833-318-0680
Enroll Online CoverMyMeds.com
Questions? Call 1-855-KESIMPTA (1-855-537-4678)

Kesimpta®
(ofatumumab) 20 mg injection

*Indicates a required field.

1 Patient Information (Please complete lab screenings specified in the Prescribing Information before initiating KESIMPTA)

First Name* Jane Last Name* Smith Email janesmith@example.com
Sex: ☐ M ☒ F Date of Birth (MM/DD/YYYY)* 09 / 16 / 1986 Home Phone (555) 555-5555 Cell Phone (555) 555-5555
Address (No PO Box) 123 Any Street
City Anytown State* NY ZIP* 10001
Caregiver First Name Jack Caregiver Last Name Smith Caregiver Email jacksmith@example.com
OK to leave voicemail on: ☒ Home Phone ☒ Cell Phone
Preferred Language: ☒ English ☐ Spanish ☐ Other:

2 Patient Authorization and Additional Consents*
I have read and agree to the Patient Authorization on page 2.
→ X *Jane Smith* Patient/Legal Guardian Signature Date of Signature (MM/DD/YYYY) 01 / 01 / 2025
*By checking this box, I agree to receive recurring marketing calls and texts from and on behalf of Novartis Pharmaceuticals Corporation. These calls and texts may be automatic or recorded in advance. The number of calls and message frequency varies. My consent is not a condition of getting any goods or services from Novartis. I can opt out of the program at any time by calling 1-855-537-4678. I can also text "STOP" to any Novartis Patient Support Ongoing Support message to opt out of texts or text "HELP" for more information about this service. Message and data rates may apply.

3 Insurance Information (Please include a copy of both sides of the insurance card)

Jane Smith Cardholder Name
First Insurance Co. (555) 555-5555 Rx Insurance Co. (555) 555-5555
Insurance Carrier Phone Number Rx Insurance Carrier Rx Phone Number
ABC1DEF12345678 12345-6789 123456 Rx BIN Number Rx PCN Number
12345-6789 ABC1DEF12345678 123456 Rx Group Number Rx ID Number

4 Provider Information
Emily Green Green Medical
First Name* Last Name* Practice Name
Address* 345 Any Avenue Nick Baker
City* Anytown State* NY ZIP* 10001 Office Contact Name
1234567890 (555) 555-5555 (555) 555-5555
NPI Number* State License # (Puerto Rico offices only) Office Contact Phone*
hi@greenmedical.com Email

5 Prescription Information
Specialty Pharmacy: Anytown Pharmacy
Preferred Specialty Pharmacy (555) 555-5555 (555) 555-5555
Phone Fax
Diagnosis Code*: ☒ ICD-10: G35 Multiple Sclerosis
Shipping Preferences: ☐ Other:
Loading Doses: ☒ Provider Address ☐ Patient Address
Pharmacy Prescription*:
KESIMPTA SensorReady® Pen
20 mg/0.4 mL single-dose prefilled pen NDC 0078-1007-68
Loading Doses:
☐ No, patient already on therapy
☒ Yes, 20 mg (0.4 mL)
SIG: 1 SC injection at week 0, 1, and 2
Qty: 3 units (No refills)
Maintenance Dose:
☒ 20 mg (0.4 mL)
SIG: 1 SC injection monthly starting at week 4
Qty: 1 SC injection, then 12 refills, or
Bridge to Commercial Coverage or Novartis Patient Assistance Foundation (NPAF):
For Bridge to Commercial Coverage or NPAF, please select applicable dose(s).
For Bridge to Commercial Coverage ONLY: Eligible patients receive KESIMPTA for free while pursuing insurance coverage. Most have commercial insurance, avoid prescription for KESIMPTA, and a denial of insurance coverage based on prior authorization request to qualify.
Loading Doses*:
☐ No, patient already on therapy
☒ Yes, 20 mg (0.4 mL)
SIG: 1 SC injection at week 0, 1, and 2
Qty: 3 units (No refills)
Maintenance Dose*:
☒ 20 mg (0.4 mL)
SIG: 1 SC injection monthly starting at week 4
Qty: 1 SC injection, then 12 refills, or

6 Provider Attestation*
Prescriber must authorize these instructions by signing at the end of this section.
I certify the above therapy is medically necessary and this information is accurate to the best of my knowledge. I certify I am the provider who has prescribed KESIMPTA to the previously identified patient and I provided the patient with a description of Alongside KESIMPTA. For the purposes of transmitting these prescriptions, I authorize NPAF, Novartis Pharmaceuticals Corporation, and its affiliates, business partners, and agents to forward as my agent, for these limited purposes, the prescriptions electronically, by facsimile, or by mail to the appropriate dispensing pharmacies. I will not attempt to seek reimbursement for free product provided to my office.
→ X *Emily Green* Provider Signature (Dispense as Written) Substitution Permissible Date of Signature (MM/DD/YYYY) 01 / 01 / 2025
ATTN: New York and Iowa providers, please submit electronic prescription to Homescripts Pharmacy, NPI #1528362076.

Complete entire form and fax to Alongside™ KESIMPTA at 1-833-318-0680
SC, subcutaneous. An incomplete Start Form may delay the start of treatment.



Give this page to your
KESIMPTA patient!

Your doctor has prescribed **KESIMPTA[®]**

It comes with membership in **Alongside[™] KESIMPTA**,
a Novartis Patient Support program.

If your doctor signed you up, here's what happens next:



We'll check your benefits

- › Expect a call from us to discuss your options,
including potential savings and product delivery



We'll mail you a welcome package

- › With some important information about
your program and quick tips for using KESIMPTA.
It should arrive in a day or two



You'll get a call from your dedicated Coordinator

- › Who has access to your membership materials,
additional training resources, and answers to any
questions about KESIMPTA and Alongside KESIMPTA.

We're in this together.



Questions? Call us.

- › **1-855-KESIMPTA (1-855-537-4678)**
8:30 AM–8:00 PM ET, Monday–Friday



Visit **www.KESIMPTA.com** for more information



*Indicates a required field.

1

Patient Information (Please complete lab screenings specified in the Prescribing Information before initiating KESIMPTA)

First Name*		Last Name*		Email	
Sex: * M F		Date of Birth (MM/DD/YYYY)*		Home Phone Cell Phone	
Address (No PO Box)					
City		State*		ZIP*	
Caregiver First Name		Caregiver Last Name		Caregiver Email	

OK to leave voicemail on: Home Phone Cell Phone

Preferred Language: English Spanish Other: _____

2

Patient Authorization and Additional Consents*

I have read and agree to the Patient Authorization on [page 2](#).

→ X

Patient/Legal Guardian Signature

Date of Signature (MM/DD/YYYY)

KESIMPTA \$0 Access Card (This box is required to access the Bridge and Copay Programs)

I have read and agree to the \$0 Access Card Terms and Conditions on page 2.

Get access to ongoing support†

I'd like to sign up for ongoing support. I'll get KESIMPTA tips, resources, and reminders from Novartis Patient Support at the mobile phone number(s) I gave above.

†By checking this box, I agree to receive recurring marketing calls and texts from and on behalf of Novartis Pharmaceuticals Corporation. These calls and texts may be automatic or recorded in advance. The number of calls and message frequency varies. My consent is not a condition of getting any goods or services from Novartis. I can opt out of the program at any time by calling 1-855-537-4678. I can also text "STOP" to any Novartis Patient Support Ongoing Support message to opt out of texts or text "HELP" for more information about this service. Message and data rates may apply.

3

Insurance Information (Please include a copy of both sides of the insurance card)

Cardholder Name		Prescription (Rx) Cardholder Name	
Insurance Carrier	Phone Number	Rx Insurance Carrier	Rx Phone Number
Cardholder ID Number	Group Number	Rx BIN Number	Rx PCN Number
		Rx Group Number	Rx ID Number

4

Provider Information

First Name*		Last Name*		Practice Name	
Address*					
City*		State*		ZIP*	
NPI Number*		State License # <small>(Puerto Rico offices only*)</small>		Email	
				Office Contact Name	
				Office Contact Phone*	
				Office Fax*	

5

Prescription Information

Specialty Pharmacy:

Preferred Specialty Pharmacy

Phone Fax

Diagnosis Code*: ICD-10: G35 Multiple Sclerosis
Other: _____

Shipping Preferences:

Loading Doses: Provider Address Patient Address

Pharmacy Prescription*:

KESIMPTA Sensoready® Pen
20 mg/0.4 mL single-dose prefilled pen NDC 0078-1007-68

Loading Doses:

No, patient already on therapy
Yes, 20 mg (0.4 mL)
SIG: 1 SC injection at week 0, 1, and 2
Qty: 3 units (No refills)

Maintenance Dose:

20 mg (0.4 mL)
SIG: 1 SC injection monthly starting at week 4
Qty: 1 SC injection, then 12 refills,
or _____ refills

Bridge to Commercial Coverage or Novartis Patient Assistance Foundation (NPAF):

For Bridge to Commercial Coverage or NPAF, please select applicable dose(s).
For Bridge to Commercial Coverage ONLY: Eligible patients receive KESIMPTA for free while pursuing insurance coverage. Must have commercial insurance, a valid prescription for KESIMPTA, and a denial of insurance coverage based on a prior authorization request to qualify.

Loading Doses*:

No, patient already on therapy
Yes, 20 mg (0.4 mL)
SIG: 1 SC injection at week 0, 1, and 2
Qty: 3 units (No refills)

Maintenance Dose*:

20 mg (0.4 mL)
SIG: 1 SC injection monthly starting at week 4
Qty: 1 SC injection, then 12 refills,
or _____ refills

6

Provider Attestation*

Prescriber must authorize these instructions by signing at the end of this section.

I certify the above therapy is medically necessary and this information is accurate to the best of my knowledge. I certify I am the provider who has prescribed KESIMPTA to the previously identified patient and I provided the patient with a description of Alongside KESIMPTA. For the purposes of transmitting these prescriptions, I authorize NPAF, Novartis Pharmaceuticals Corporation, and its affiliates, business partners, and agents to forward as my agent, for these limited purposes, the prescriptions electronically, by facsimile, or by mail to the appropriate dispensing pharmacies. I will not attempt to seek reimbursement for free product provided to my office.

→ X

Provider Signature (Dispense as Written)

Substitution Permissible

Date of Signature (MM/DD/YYYY)

ATTN: New York and Iowa providers, please submit electronic prescription to Homescripts Pharmacy, NPI #1528362076.

Complete entire form and fax to Alongside™ KESIMPTA at **1-833-318-0680**

An incomplete Start Form may delay the start of treatment.

Patient Authorization. I authorize my health care professionals, pharmacies and health insurers, and their service providers (“Providers”) to disclose information relating to my insurance benefits, medical condition, treatment and prescription details (“Personal Information”) to Novartis Pharmaceuticals Corporation, its affiliates and service providers (“Novartis”) and the Novartis Patient Assistance Foundation, Inc., and its service providers (“NPAF”) so they can provide the following support services (the “Services”):

- Help coordinate insurance coverage for, access to, and receipt of my medication.
- Communicate with me about possible financial assistance, including Novartis copay or NPAF programs, and, if I am enrolled, administer my participation in those programs.
- Communicate with me about my medication and treatment, including reminders, health and lifestyle tips, and product and other related information. Communications may be customized based on Personal Information obtained from my Providers.
- Conduct quality assurance and other internal business activities and ask for feedback related to the Services or my treatment.

In delivering the Services, Novartis and NPAF may share my Personal Information with each other, with my Providers, or with government agencies or other financial assistance programs that might help me pay for my medication. They may combine information collected from me with information collected from other sources and use that information to administer the Services. My pharmacies or other health care professionals may receive payment from Novartis or NPAF for providing certain Services, such as medication or refill reminders, based on my enrollment or participation. Once I authorize disclosure of my Personal Information, it may no longer be protected by federal health privacy law and applicable state laws.

I understand I do not have to sign this Authorization to get my medication or insurance coverage, that I have a right to a copy, and can cancel this Authorization at any time by calling 1-855-537-4678 or by writing to:

PO Box 2971
850 Twin Rivers Dr
Columbus, OH 43216-9532

OR

Customer Interaction Center
Novartis Pharmaceuticals Corporation
One Health Plaza
East Hanover, NJ 07936-1080

This Authorization will expire 5 years after I sign it, or earlier if required by state law, unless I cancel it sooner. If I cancel it, I may no longer qualify for Services from Novartis or NPAF, but it will not impact my Provider’s treatment or my insurance benefits. I also understand that if a Provider is disclosing my Personal Information to Novartis or NPAF on an authorized, ongoing basis, my cancellation will be effective with respect to that Provider as soon as they receive notice of my cancellation. Cancellation will not affect prior uses or disclosures.

\$0 Access Card Terms and Conditions

Limitations apply. Offer not valid under Medicare, Medicaid, or any other federal or state health insurance program. Patients with commercial insurance who are initially denied coverage may receive free KESIMPTA® (ofatumumab) for up to 12 months while seeking coverage. Patients with commercial insurance who have coverage for KESIMPTA may receive up to \$18,000 in annual copay benefits. Novartis reserves the right to rescind, revoke, or amend this program without notice. Additional limitations may apply. See complete Terms & Conditions at start.kesimpta.com.

Bridge Program: Must have commercial insurance, a valid prescription for KESIMPTA, and a denial of insurance coverage based on a prior authorization requirement to qualify. Eligible patients may receive a monthly maintenance dose for up to 12 months or until insurance coverage approval, whichever occurs first. Not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, VA, DoD or any other federal or state program, or where prohibited by law. No purchase necessary. Program is not health insurance, nor is participation a guarantee of insurance coverage. Other limitations may apply. Novartis reserves the right to rescind, revoke, or amend this Program without notice.

IMPORTANT SAFETY INFORMATION (CONT)

Hepatitis B Virus: Reactivation: No reports of HBV reactivation in patients with MS treated with KESIMPTA. However, HBV reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, has occurred in patients treated with ofatumumab at higher intravenous doses for chronic lymphocytic leukemia (CLL) than the recommended dose in MS and in patients treated with other anti-CD20 antibodies.

Infection: KESIMPTA is contraindicated in patients with active hepatitis B disease. Fatal infections caused by HBV in patients who have not been previously infected have occurred in patients treated with ofatumumab at higher intravenous doses for CLL than the recommended dose in MS. Perform HBV screening in all patients before initiation of KESIMPTA. Patients who are negative for HBsAg and positive for HB core antibody [HBcAb+] or are carriers of HBV [HBsAg+], should consult liver disease experts before starting and during KESIMPTA treatment.

Progressive Multifocal Leukoencephalopathy: No cases of progressive multifocal leukoencephalopathy (PML) have been reported for KESIMPTA in RMS clinical studies; however, PML resulting in death has occurred in patients being treated with ofatumumab at higher intravenous doses for CLL than the recommended dose in MS. In addition, JC virus infection resulting in PML has also been observed in patients treated with other anti-CD20 antibodies and other MS therapies. If PML is suspected, withhold KESIMPTA and perform an appropriate diagnostic evaluation. If PML is confirmed, KESIMPTA should be discontinued.

Vaccinations: Administer all immunizations according to immunization guidelines: for live or live-attenuated vaccines at least 4 weeks and, whenever possible at least 2 weeks prior to starting KESIMPTA for inactivated vaccines. The safety of immunization with live or live-attenuated vaccines following KESIMPTA therapy has not been studied. Vaccination with live or live-attenuated vaccines is not recommended during treatment and after discontinuation until B-cell repletion.

Vaccination of Infants Born to Mothers Treated with KESIMPTA During Pregnancy. For infants whose mother was treated with KESIMPTA during pregnancy, assess B-cell counts prior to administration of live or live-attenuated vaccines. If the B-cell count has not recovered in the infant, do not administer the vaccine as having depleted B-cells may pose an increased risk in these infants.

Injection-Related Reactions and Hypersensitivity Reactions: KESIMPTA can result in systemic injection-related reactions and hypersensitivity reactions, which may be serious or life-threatening. Injection-related reactions with systemic symptoms occurred most commonly within 24 hours of the first injection, but were also observed with later injections. There were no life-threatening injection reactions in RMS clinical studies.

In the post-marketing setting, additional systemic injection-related reactions and hypersensitivity reactions have been reported, including anaphylaxis, angioedema, pruritus, rash, urticaria, erythema, bronchospasm, throat irritation, oropharyngeal pain, dyspnea, pharyngeal or laryngeal edema, flushing, hypotension, dizziness, nausea, and tachycardia. Most cases were not serious and occurred with the first injection. Symptoms of systemic injection-related reactions may be clinically indistinguishable from acute hypersensitivity reactions.

The first injection of KESIMPTA should be performed under the guidance of an appropriately trained health care professional. If systemic injection-related reactions occur, initiate appropriate therapy. Patients who experience symptoms of systemic injection-related reactions or hypersensitivity reactions with KESIMPTA should be instructed to seek immediate medical attention. If local injection-related reactions occur, symptomatic treatment is recommended.

Reduction in Immunoglobulins: As expected with any B-cell depleting therapy, decreased immunoglobulin levels were observed. Monitor the levels of quantitative serum immunoglobulins during treatment, especially in patients with opportunistic or recurrent infections and after discontinuation of therapy until B-cell repletion. Consider discontinuing KESIMPTA therapy if a patient with low immunoglobulins develops a serious opportunistic infection or recurrent infections, or if prolonged hypogammaglobulinemia requires treatment with intravenous immunoglobulins.

Fetal Risk: Based on animal data, KESIMPTA can cause fetal harm due to B-cell lymphopenia and reduce antibody response in offspring exposed to KESIMPTA in utero. Transient peripheral B-cell depletion and lymphocytopenia have been reported in infants born to mothers exposed to other anti-CD20 B-cell depleting antibodies during pregnancy. Advise females of reproductive potential to use effective contraception while receiving KESIMPTA and for at least 6 months after the last dose.

Most common adverse reactions (>10%) are upper respiratory tract infection, headache, injection-related reactions, and local injection-site reactions.

Please see additional Important Safety Information on [the front cover](#).

Please see full Prescribing Information, including Medication Guide [here](#).

